

# Global Barriers to Eye Care Access: A Systematic Review and Meta-Analysis of Gender, Geographic, and Economic Disparities



Debanjali Bhattacharjee<sup>1</sup>, Ankit Sanjay Varshney<sup>2</sup>, Neelam Dhankar<sup>3</sup>

<sup>1,3</sup>SHS, Sushant University, Gurgaon, India

<sup>2</sup>Shree Bharatimaiya College of Optometry, Surat, India

## ABSTRACT

Access to eye care remains a major global challenge, particularly in low- and middle-income countries and rural areas, where financial, geographic, and awareness-related barriers contribute to preventable blindness. This systematic review and meta-analysis synthesized 45 studies from 28 countries including 1.2 million adults, following PRISMA 2020 guidelines. The pooled prevalence of financial, geographic, and awareness barriers was 58.7%, 41.2%, and 39.8%, respectively, with women, rural residents, and low-income groups disproportionately affected. Subgroup analyses confirmed significantly higher odds of access barriers in low-resource settings. Cataract and uncorrected refractive errors remained the leading global causes of blindness. Findings highlight the need for equity-focused interventions such as universal health coverage, telemedicine, mobile clinics, and community-based education. Addressing these systemic barriers is essential to achieve Vision 2030 goals and reduce avoidable blindness worldwide.

**Key-words:** Health Services Accessibility, Blindness, Developing Countries, Rural Health, Rural Health Services.

**How to Cite this Article:** Bhattacharjee D, Varshney AS, Dhankar N. Global Barriers to Eye Care Access: A Systematic Review and Meta-Analysis of Gender, Geographic, and Economic Disparities. 2026;42(2):220-228. Doi:10.36351/pjo.v42i2.2132

---

*Correspondence: Debanjali Bhattacharjee  
SHS, Sushant University, Gurgaon, India  
Email: optomdebanjali@gmail.com*

---

*Received: June 25, 2025  
Revised: March 1, 2026  
Accepted: March 15, 2026*

## INTRODUCTION

Vision impairment and blindness remain major global public health challenges, affecting more than 2.2 billion people worldwide, with at least 1 billion cases considered preventable or treatable.<sup>1,2</sup> Despite advances in ophthalmic diagnostics, surgical techniques, and global eye-health initiatives, access to eye-care services remains highly unequal across populations.<sup>3</sup> Disparities are particularly evident in low- and middle-income countries (LMICs) and rural regions where healthcare infrastructure, trained

personnel, and financial protection mechanisms are limited.<sup>3,4</sup> These inequalities contribute not only to preventable vision loss but also to substantial economic and social consequences, as visual impairment reduces productivity, limits employment opportunities, and increases healthcare expenditures.<sup>4,5</sup>

Globally, cataracts and uncorrected refractive errors remain the leading causes of vision impairment and blindness, despite being largely treatable through cost-effective interventions such as cataract surgery and refractive correction.<sup>6,7</sup> However, the uptake of these services continues to be hindered by multiple barriers, including financial constraints, geographic inaccessibility, limited health infrastructure, and lack of awareness of available services.<sup>8,9</sup> These challenges disproportionately affect vulnerable populations such as women, older adults, individuals with low socioeconomic status, and residents of remote communities, thereby reinforcing existing health inequalities.<sup>10,11</sup>

Recognizing the growing burden of avoidable blindness, global initiatives such as Vision 2020, the Lancet Global Health Commission on Global Eye Health, and the Sustainable Development Goals (SDGs) emphasize the importance of achieving universal and equitable access to eye-care services.<sup>12,13</sup> These frameworks advocate strengthening primary eye-care systems, expanding community-based screening programs, and adopting innovative service delivery models such as teleophthalmology and mobile eye-care services.<sup>14</sup>

Despite these initiatives, many existing studies examining barriers to eye-care access remain region-specific or focused on individual determinants, limiting a comprehensive understanding of global disparities. Therefore, this systematic review and meta-analysis aim to synthesize global evidence on financial, geographic, and awareness-related barriers to eye-care access and quantify disparities across different populations and settings.

## METHODS

This systematic review and meta-analysis followed PRISMA 2020 guidelines to ensure methodological transparency and reporting quality. Although the review protocol was not registered in PROSPERO, the study methodology, eligibility criteria, and analytic plan were defined a priori to maintain methodological rigor.

PubMed, Scopus, and Google Scholar were searched for studies published between January 2000 and March 2024 using MeSH and free-text terms (e.g., *barriers to eye care*, *blindness causes*, *cataract surgery access*). Filters restricted results to English-language, peer-reviewed studies of adults ( $\geq 18$  years). Grey literature from WHO, government health reports, and NGO data was screened, and duplicates were removed with Zotero.

Studies were included if they reported financial, geographic, or awareness-related barriers to eye care in adults. Both quantitative and qualitative designs were eligible if methodologically sound. Paediatric studies, non-empirical work, studies without outcome data, and retracted or duplicate articles were excluded.

Two reviewers independently extracted study characteristics, demographics, barrier types, and statistical outputs. Discrepancies were resolved by consensus, with a third reviewer as needed. Study

quality was assessed using the Newcastle-Ottawa Scale for observational studies, CASP for qualitative work, and RoB 2/ROBINS-I for trials.

A random-effects meta-analysis estimated pooled prevalence and odds ratios. Heterogeneity was quantified using the  $I^2$  statistic, and subgroup analyses examined differences by geography, income, gender, and age. Meta-regression tested covariate effects. Analyses were performed in RevMan 5.4 and R (packages *meta* and *metaphor*). Publication bias was assessed with funnel plots, Egger's test, and trim-and-fill methods.

Only publicly available data were analysed; thus, ethical approval was not required. All included studies were reviewed for compliance with ethical standards, including informed consent.

## RESULTS

This review included 45 studies from 28 countries, encompassing approximately 1.2 million participants (Figure 1). Reported barriers to eye care were categorised as financial, geographic, and awareness related. Key study characteristics, including study design, population demographics, geographic setting, and identified barriers, are summarised and provided as a supplementary file (Appendix A)

**Financial Barriers:** This was the most common barrier, with a pooled prevalence of 58.7% (95% CI: 54.2–63.1%) (Figure 2). Rates were higher in LMICs (65.4%) than High Income Countries (HICs) with 42.1% (OR = 2.2, 95% CI: 1.8–2.7), and in rural (69.2%) vs. urban populations (47.5%) (OR = 2.3, 95% CI: 1.9–2.8). Women reported higher prevalence (62.3%) than men (55.1%) (OR = 1.4, 95% CI: 1.1–1.7).

**Geographic Barriers:** Pooled prevalence was 41.2% (95% CI: 36.8–45.7%), significantly higher in LMICs (48.9%) than HICs (28.3%) (OR = 2.1, 95% CI: 1.7–2.6). Rural residents were disproportionately affected (56.1% vs. 32.4% urban; OR = 2.6), as were women (44.5% vs. 37.8%; OR = 1.3) (Figure 3).

**Awareness Barriers:** Overall prevalence was 39.8% (95% CI: 35.4–44.3%), with higher rates in LMICs (45.6%) vs. HICs (28.7%) (OR = 2.0), and in rural (51.2%) vs. urban (29.6%) settings (OR = 2.5). Women reported slightly greater barriers than men (42.7% vs. 36.9%; OR = 1.2).

**Combined Barriers:** Financial and geographic barriers overlapped in 62.4% of participants, rising to 69.8% in LMICs and 74.5% in rural areas (ORs = 2.3–2.8),(Figure 4).

**Subgroup Trends:** Across domains, women, rural residents, and low-income groups consistently experienced greater barriers ( $p < 0.05$ ), confirming intersectional disadvantage (Figure 5).

**Global Causes of Blindness:** Cataracts accounted for 51% of global cases, followed by uncorrected refractive errors (20%), glaucoma (8%), diabetic retinopathy (4.8%), and AMD (5%) (Figure 6). Trends from 2010–2024 showed persistent cataract burden and increasing diabetic retinopathy (Figure 7).

**Interventions:** Strategies such as telemedicine, subsidized healthcare, mobile clinics, and school-based screening demonstrated potential to reduce barriers (Figures 8, 9).

**Statistical Analysis:** Heterogeneity was high ( $I^2 > 75\%$ ), partly explained by economic status in meta-regression. Sensitivity analyses supported robustness. Funnel plots suggested modest publication bias for financial barriers, but Egger’s test was nonsignificant. Subgroup odds ratios confirmed disparities by gender, geography, and income (Figure 10).

## DISCUSSION

This systematic review and meta-analysis consolidates global evidence on barriers to eye care and aligns with findings from major global eye-health initiatives and epidemiological analyses of vision impairment.<sup>12,13,15,16</sup> The pooled prevalence of financial (58.7%), geographic (41.2%), and awareness-related barriers (39.8%) indicates that despite major advances in ophthalmic technologies, service delivery models, and global initiatives, large segments of the population remain unable to access essential eye-care services.<sup>1,2</sup> These findings are concerning because most causes of vision impairment are preventable or treatable with timely interventions.

Subgroup analyses further highlight the unequal distribution of barriers across populations. Individuals living in LMICs and rural settings experience the highest burden of access limitations. Financial barriers affect nearly two-thirds of individuals in LMICs and rural communities, compared with substantially lower rates in HICs and urban areas.<sup>3,4</sup> These disparities reflect systemic weaknesses in health-financing systems, limited insurance coverage, and high out-of-pocket healthcare expenditures.<sup>17</sup> Geographic barriers also remain substantial, particularly in regions where




Figure 1: PRISMA 2020 flow diagram of study selection.




Figure 2: Forest Plot of Pooled Prevalence Estimates and Odds Ratios (with 95% CI) for Eye Care Access Barriers, Stratified by Rural/Urban and LMIC/HIC Status.





Figure 3: Bar Charts Illustrating the Prevalence of Financial, Geographic, and Awareness-Related Barriers to Eye Care Access: Regional (LMICs vs. HICs), Geographic (Urban vs. Rural), and Gender (Women vs. Men) Comparisons.

ophthalmic infrastructure and trained personnel are concentrated in urban centres.<sup>6,18</sup>


Gender inequities further compound these access challenges. Women consistently report greater financial, geographic, and awareness-related barriers to eye care utilization.<sup>10,11</sup> This may be attributed to intersecting socioeconomic and cultural factors such as lower financial independence, mobility restrictions,

caregiving responsibilities, and limited participation in healthcare decision-making. Evidence from broader healthcare research suggests that women with limited socioeconomic autonomy often face systemic obstacles in accessing healthcare services.<sup>19</sup>

The continued dominance of cataract and uncorrected refractive errors as leading causes of blindness worldwide also reflects gaps in delivering



**Figure 4:** Heatmap Illustrating the Combined Prevalence of Financial, Geographic, and Awareness-Related Barriers to Eye Care Access: LMICs vs. HICs and Urban vs. Rural Comparisons.



**Figure 5:** Boxplot of Barrier Prevalence Across Gender, Socioeconomic Status (LMICs/HICs), and Geographic Setting (Rural/Urban)

cost-effective eye-care interventions.<sup>5,6</sup> Cataract surgery and refractive correction are among the most cost-effective health interventions globally, yet their utilization remains suboptimal due to the barriers identified in this review.<sup>9,20</sup> In contrast, HICs are experiencing an epidemiological shift toward chronic retinal diseases such as age-related macular degeneration and diabetic retinopathy, largely driven by ageing populations and the increasing prevalence of non-communicable diseases.<sup>16,21</sup>

The substantial heterogeneity observed across studies ( $I^2 > 75\%$ ) suggests that barriers to eye care are influenced by complex contextual factors, including health-system capacity, geographic accessibility,

population demographics, and sociocultural determinants. Therefore, strategies designed to improve eye-care access must be tailored to local contexts rather than relying on universal global approaches.


### Policy and Practice Implications

In line with global strategies such as Vision 2030 and the Sustainable Development Goals,<sup>12-14</sup> a multipronged approach is required to address the structural determinants of eye-care access.


Financial barriers can be reduced by integrating eye-care services within universal health coverage frameworks and expanding financial protection mechanisms for vulnerable populations.<sup>17,20</sup> Cross-subsidization models and public-private partnerships may further expand access to affordable optical and surgical services.<sup>22</sup>

Geographic barriers require strengthening rural healthcare infrastructure through mobile outreach programs, satellite clinics, and decentralized vision centres.<sup>6,18</sup> Teleophthalmology also represents a promising strategy for extending specialist services to remote communities.<sup>14</sup>

Awareness barriers may be addressed through culturally appropriate community education programs, school-based vision screening initiatives, and digital health campaigns aimed at increasing awareness of preventable eye diseases.<sup>8</sup>



**Figure 6:** Proportional Distribution of Global Causes of Blindness: Cataracts, Refractive Errors, Glaucoma, Diabetic Retinopathy, AMD, and Other Causes.



**Figure 7:** Line Graph Showing the Prevalence of Cataracts, Uncorrected Refractive Errors, Glaucoma, and Diabetic Retinopathy from 2010 to 2024.

Addressing gender and age disparities is equally important. Targeted policies that reduce women’s financial and mobility constraints, expand elder-friendly eye-care services, and engage community organizations can help improve equitable access to care.

## CONCLUSION

This study highlights financial, geographic, and awareness-related barriers as critical impediments to

achieving equitable access to eye care services globally. Women, rural populations, and individuals in LMICs bear the highest burden of these barriers. Addressing these challenges requires integrated, context-specific interventions that leverage universal health coverage, telemedicine technologies, and targeted health education campaigns.

These findings align with global initiatives like Vision 2030 and the Sustainable Development Goals, offering a clear roadmap for policymakers, healthcare providers, and international stakeholders to implement sustainable and equitable solutions for preventable blindness and vision impairment worldwide.


**Funding:** This study was not funded by any organization.

**Patient’s Consent:** Researchers followed the guidelines set forth in the Declaration of Helsinki.


**Conflict of Interest:** Authors declared no conflict of interest.

## ACKNOWLEDGEMENTS


The authors sincerely acknowledge the contributions of all researchers whose primary studies were included in this review. We thank our institutions for their academic support and access to library resources. We also appreciate the assistance provided by our head of department and staff members during data organization and formatting.



**Figure 8:** Radar Chart and Sankey Diagram Comparing the Impact of Telemedicine and Subsidized Healthcare on Reducing Financial, Geographic, and Awareness Barriers to Eye Care Access.



**Figure 9:** Color-Coded Comparison of Interventions: Effectiveness and Implementation Challenges in Eye Care Access Programs.



**Figure 10:** Forest Plot and Funnel Plot Assessing the Impact of Financial Barriers to Eye Care Access Across Gender, Geographic, and Income Subgroups.

## REFERENCES

- Vashist P, Senjam SS, Gupta V, Gupta N, Shamanna BR, Wadhvani M, et al.** Blindness and visual impairment and their causes in India: Results of a nationally representative survey. *PLoS One*. 2022;**17(7)**:e0271736. Doi: 10.1371/journal.pone.0271736.
- Yekta A, Hooshmand E, Saatchi M, Ostadimoghaddam H, Asharlous A, Taheri A, et al.** Global Prevalence and Causes of Visual Impairment and Blindness in Children: A Systematic Review and Meta-Analysis. *J Curr Ophthalmol*. 2022;**34(1)**:1-15. Doi: 10.4103/joco.joco\_135\_21.
- Seidu AA, Agbadi P, Duodu PA, Dey NEY, Duah HO, Ahinkorah BO.** Prevalence and sociodemographic factors associated with vision difficulties in Ghana, Gambia, and Togo: a multi-country analysis of recent multiple Indicator cluster surveys. *BMC Public Health*. 2021;**21(1)**:2148. Doi: 10.1186/s12889-021-12193-7.
- Marques AP, Ramke J, Cairns J, Butt T, Zhang JH, Muirhead D, et al.** Global economic productivity losses from vision impairment and blindness. *EclinMed*. 2021;**35**:100852. Doi: 10.1016/j.eclinm.2021.100852.
- Smith TS, Frick KD, Holden BA, Fricke TR, Naidoo KS.** Potential lost productivity resulting from the global burden of uncorrected refractive error. *Bull World Health Organ*. 2009;**87(6)**:431-437. Doi: 10.2471/blt.08.055673.
- Correia M, Das T, Magno J, Pereira BM, Andrade V, Limburg H, et al.** Prevalence and causes of blindness, visual impairment, and cataract surgery in Timor-Leste. *Clin Ophthalmol*. 2017;**11**:2125-2131. Doi: 10.2147/OPTH.S146901.
- Oye JE, Kuper H.** Prevalence and causes of blindness and visual impairment in Limbe urban area, South West Province, Cameroon. *Br J Ophthalmol*. 2007;**91(11)**:1435-1439. Doi: 10.1136/bjo.2007.115840.
- Elam AR, Lee PP.** Barriers to and Suggestions on Improving Utilization of Eye Care in High-Risk Individuals: Focus Group Results. *Int Sch Res Notices*. 2014;**2014**:527831. Doi: 10.1155/2014/527831.
- Marmamula S, Khanna RC, Shekhar K, Rao GN.** A population-based cross-sectional study of barriers to uptake of eye care services in South India: the Rapid Assessment of Visual Impairment (RAVI) project. *BMJ Open*. 2014;**4(6)**:e005125. Doi: 10.1136/bmjopen-2014-005125.
- Neyhouser C, Quinn I, Hillgrove T, Chan R, Chhea C, Peou S, et al.** A qualitative study on gender barriers to eye care access in Cambodia. *BMC Ophthalmol*. 2018;**18(1)**:217. Doi: 10.1186/s12886-018-0890-3.
- Sabherwal S, Sood I, Garg GK, DasGupta S, Nagappan S, Reddy PA, et al.** Gender inequity in eyecare: variation by service level and location in North India. *Indian J Public Health Res Dev*. 2020;**11(7)**:784-7990. Doi:10.37506/ijphrd.v11i7.10183

12. **Zhang JH, Ramke J, Jan C, Bascaran C, Mwangi N, Furtado JM, et al.** Advancing the Sustainable Development Goals through improving eye health: a scoping review. *Lancet Planet Health.* 2022;**6(3)**:e270-e280. Doi: 10.1016/S2542-5196(21)00351-X.
13. **Burton MJ, Ramke J, Marques AP, Bourne RRA, Congdon N, Jones I, et al.** The Lancet Global Health Commission on Global Eye Health: vision beyond 2020. *Lancet Glob Health.* 2021;**9(4)**:e489-e551. Doi: 10.1016/S2214-109X(20)30488-5.
14. **Prathiba V, Rema M.** Teleophthalmology: a model for eye care delivery in rural and underserved areas of India. *Int J Family Med.* 2011;**2011**:683267. Doi: 10.1155/2011/683267.
15. **Ramke J, Evans JR, Habtamu E, Mwangi N, Silva JC, Swenor BK, et al.** Grand Challenges in Global Eye Health study group. Grand Challenges in global eye health: a global prioritisation process using Delphi method. *Lancet Healthy Longev.* 2022;**3(1)**:e31-e41. Doi: 10.1016/S2666-7568(21)00302-0.
16. **Flaxman SR, Bourne RRA, Resnikoff S, Ackland P, Braithwaite T, Cicinelli MV, et al.** Vision Loss Expert Group of the Global Burden of Disease Study. Global causes of blindness and distance vision impairment 1990-2020: a systematic review and meta-analysis. *Lancet Glob Health.* 2017;**5(12)**:e1221-e1234. Doi: 10.1016/S2214-109X(17)30393-5.
17. **Wittenborn JS, Zhang X, Feagan CW, Crouse WL, Shrestha S, Kemper AR, et al.** Vision Cost-Effectiveness Study Group. The economic burden of vision loss and eye disorders among the United States population younger than 40 years. *Ophthalmology.* 2013;**120(9)**:1728-1735. Doi: 10.1016/j.opht.2013.01.068.
18. **Owusu-Afriyie B, Peter N, Ivihi F, Kopil I, Gende T.** Barriers to the uptake of eye care services: A cross-sectional survey from rural and urban communities. *PLoS One.* 2024;**19(8)**:e0308294. Doi: 10.1371/journal.pone.0308294.
19. **Sharts-Hopko NC, Smeltzer S, Ott BB, Zimmerman V, Duffin J.** Healthcare experiences of women with visual impairment. *Clin Nurse Spec.* 2010;**24(3)**:149-153. Doi: 10.1097/NUR.0b013e3181d82b89.
20. **Congdon N, O'Colmain B, Klaver CC, Klein R, Muñoz B, Friedman DS, et al.** Eye Diseases Prevalence Research Group. Causes and prevalence of visual impairment among adults in the United States. *Arch Ophthalmol.* 2004;**122(4)**:477-85. Doi: 10.1001/archoph.122.4.477.
21. **Zaback T, Lam S, Randall J, Field T, Brinks MV.** Access to Eye Care Before and After Vision Loss: A Qualitative Study Investigating Eye Care Among Persons Who Have Become Blind. *Qual Rep.* 2020;**25(6)**:1473-1488. PMID: 32924032.
22. **Kovai V, Rao GN, Holden B, Sannapaneni K, Bhattacharya SK, Khanna R.** Comparison of patient satisfaction with services of vision centers in rural areas of Andhra Pradesh, India. *Indian J Ophthalmol.* 2010;**58(5)**:407-413. Doi: 10.4103/0301-4738.67056.

### **Authors Designation and Contribution**

Debanjali Bhattacharjee; Assistant Professor:  
*Concepts, Literature search, Data acquisition, Statistical analysis, Manuscript preparation, Manuscript editing.*

Ankit Sanjay Varshney; Associate Professor:  
*Concepts, Design, Data analysis, Manuscript preparation, Manuscript review.*

